

**§ 556.580 Robenidine hydrochloride.**

Tolerances are established for residues of robenidine hydrochloride in edible tissues of chickens as follows:

(a) 0.2 part per million in skin and fat.

(b) 0.1 part per million (negligible residue) in edible tissues other than skin and fat.

**§ 556.590 Salicylic acid.**

A tolerance of zero is established for residues of salicylic acid in milk from dairy animals.

**§ 556.594 Sarafloxacin.**

A tolerance for residues of sarafloxacin in edible turkey and broiler chickens tissues is not required.

[60 FR 50098, Sept. 28, 1995]

**§ 556.600 Spectinomycin.**

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of spectinomycin is 25 micrograms per kilogram of body weight per day.

(b) *Chickens and turkeys*. A tolerance of 0.1 part per million (ppm) for negligible residues of spectinomycin in uncooked edible tissues of chickens and turkeys is established.

(c) *Cattle*. A tolerance of 4 ppm for parent spectinomycin (marker residue) in kidney (target tissue) is established. A tolerance of 0.25 ppm for parent spectinomycin in cattle muscle is established.

[63 FR 24107, May 1, 1998; 63 FR 38304, July 16, 1998]

**§ 556.610 Streptomycin.**

Tolerances are established for residues of streptomycin in uncooked, edible tissues of chickens, swine, and calves of 2.0 parts per million (ppm) in kidney and 0.5 ppm in other tissues.

[58 FR 47211, Sept. 8, 1993]

**§ 556.620 Sulfabromomethazine sodium.**

Tolerances for residues of sulfabromomethazine sodium in food are established as follows:

(a) In the uncooked edible tissues of cattle at 0.1 part per million (negligible residue).

(b) In milk at 0.01 part per million (negligible residue).

[47 FR 30244, July 13, 1982]

**§ 556.625 Sodium sulfachloropyrazine monohydrate.**

A tolerance of zero is established for residues of sodium sulfachloropyrazine monohydrate in the uncooked edible tissues of chickens.

**§ 556.630 Sulfachlorpyridazine.**

A tolerance of 0.1 part per million is established for negligible residues of sulfachlorpyridazine in uncooked edible tissues of calves and swine.

**§ 556.640 Sulfadimethoxine.**

Tolerances are established for residues of sulfadimethoxine in edible products of animals as follows:

(a) In the uncooked edible tissues of chickens, turkeys, cattle, ducks, salmonids, and catfish at 0.1 part per million (negligible residue).

(b) In milk at 0.01 part per million (negligible residue).

[40 FR 13942, Mar. 27, 1975, as amended at 49 FR 46371, Nov. 26, 1984; 51 FR 18884, May 23, 1986]

**§ 556.650 Sulfaethoxypyridazine.**

Tolerances for residues of sulfaethoxypyridazine in food are established as follows:

(a) Zero in the uncooked edible tissues of swine and in milk.

(b) 0.1 part per million (negligible residue) in uncooked edible tissues of cattle.

**§ 556.660 Sulfamerazine.**

A tolerance of zero is established for residues of sulfamerazine (N<sup>1</sup>-[4-methyl-2-pyrimidinyl]sulfanilamide) in the uncooked edible tissues of trout.

**§ 556.670 Sulfamethazine.**

A tolerance of 0.1 part per million is established for negligible residues of sulfamethazine in the uncooked edible tissues of chickens, turkeys, cattle, and swine.

[47 FR 25323, June 11, 1982]

**§ 556.680 Sulfanitran.**

A tolerance of zero is established for residues of sulfanitran (acetyl(*p*-nitro-